

Heat transport and spin-charge separation in the normal state of high temperature superconductors

Qimiao Si

Department of Physics and Astronomy, Rice University, Houston, TX 77251-1892

Hill et al. have recently measured both the thermal and charge conductivities in the normal state of a high temperature superconductor. Based on the vanishing of the Wiedemann-Franz ratio in the extrapolated zero temperature limit, they conclude that the charge carriers in this material are not fermionic. Here I make a simple observation that the prefactor in the temperature dependence of the measured thermal conductivity is unusually large, corresponding to an extremely small energy scale $T_0 \approx 0.15$ K. I argue that T_0 should be interpreted as a collective scale. Based on model-independent considerations, I also argue that the experiment leads to two possibilities: 1) The charge-carrying excitations are non-fermionic. And much of the heat current is in fact carried by distinctive charge-neutral excitations; 2) The charge-carrying excitations are fermionic, but a subtle ordering transition occurs at T_0 .

Spin-charge separation has long been proposed to describe the normal state of the high temperature superconductors¹, and continues to be the subject of extensive current work. It has, however, received no unambiguous experimental support, many indirect evidences notwithstanding². One natural means to probe spin-charge separation is to compare the temperature dependences of spin transport and charge transport properties³⁻⁵. Spin injection experiments have recently been carried out in the cuprates⁶⁻⁸; these experiments, however, have yet to yield quantitative information about spin transport. A less direct alternative to spin transport in this context is heat transport, if the electronic contribution can be unambiguously separated from the phononic one. While such a separation is easy to achieve in highly conductive metals, it is in general very difficult for strongly correlated metals⁹.

Very recently, Hill et al.¹⁰ have extracted the electronic contribution to the thermal conductivity (κ) in the normal state of an optimally electron-doped PCCO. This is achieved at very low temperatures, where scattering is dominated by elastic processes. The normal state arises in a magnetic field applied along the c -axis $H \approx 13$ T, which is above the bulk upper critical field H_{c2} . (The thermal conductivity is field-independent at $H > H_{c2}$.) At roughly the same magnetic field, the electrical conductivity (σ) is essentially temperature-independent, up to small weak-localization-like corrections. The authors take advantage of the fact that the measured zero-field thermal conductivity $\kappa(H=0)$ goes to zero in approximately a T^3 fashion, in sharp contrast to what is generally expected for a disordered d-wave superconductor¹¹. They assume that $\kappa(H=0)$ is entirely due to phonons, and identify the electronic contribution to the normal-state thermal conductivity as $\kappa_e = \kappa(H \approx 13 \text{ T}) - \kappa(H=0)$. κ_e is found to have an asymptotic low temperature form

$$\lim_{T \rightarrow 0} \kappa_e \sim T^{\alpha+1} \quad (1)$$

where $\alpha \approx 2.6$. This is in strong violation of the Wiedemann-Franz (WF) law: According to this law, α should be equal to zero reflecting the linear temperature dependence of the specific heat of the fermionic quasiparticles in a Fermi liquid. From this perspective, the experiment implies a missing entropy. The authors conclude that the carriers of the charge current are not fermionic.

Here I point out that the prefactor in the temperature dependence of the thermal conductivity should be considered to be unusually large, if the dominant contribution to κ_e were due to non-fermionic charge-carrying excitations. To see this, we note that the measured κ_e becomes larger than

$$\kappa_{WF} \equiv L_0 T \sigma \quad (2)$$

at a temperature $T_0 \approx 0.15$ K. (Here $L_0 \equiv \pi^2 k_B^2 / 3e^2$ is the Lorenz number of a Fermi liquid.) Namely, the experimental data can be cast in the form,

$$\lim_{T \rightarrow 0} \frac{\kappa_e}{\kappa_{WF}} \approx \left(\frac{T}{T_0} \right)^\alpha \quad (3)$$

Using $\sigma = e^2 (dn/d\mu) D_{charge}$, where $dn/d\mu$ and D_{charge} are the electronic compressibility and charge diffusion constant, respectively, we can rewrite

$$\kappa_{WF} = L_0 T (e^2/m) n_{charge} \tau_0 \frac{(dn/d\mu)}{N_0} \frac{D_{charge}}{D_0} \quad (4)$$

where the subscript 0 labels quantities in the absence of interactions. It follows that κ_{WF} does not explicitly depend on the velocity and, equivalently, the bandwidth, of the charge-carrying excitations.

If the measured κ_e is dominated by the contribution of the charge-carrying excitations, we can express κ_e/κ_{WF} in the following general form:

$$\frac{\kappa_e}{\kappa_{WF}} = \frac{C_v}{C_v^0} \frac{N_0}{(dn/d\mu)} \frac{D_{heat}}{D_{charge}} \quad (5)$$




FIG. 1. Schematics of two possible pictures. The solid lines give the ratio of the total electronic thermal conductivity, κ_e , to κ_{WF} , defined in Eq. (2). In a), the dotted line describes the contribution from charge-carrying excitations; the remaining contribution comes from charge-neutral excitations. In b), the excitations are fermionic and the low temperature behavior reflects an ordering transition around T_0 .

where D_{heat} is the entropy diffusion constant and C_v the specific heat due to the charge-carrying excitations. In this case, the scattering times for both the heat and charge currents reflect the elastic scattering time of the same excitation. As a result, the ratio D_{heat}/D_{charge} is expected to depend only on equilibrium interaction parameters¹². These interaction parameters should mostly cancel out in the product $\frac{D_{heat} N_0}{D_{charge} (dn/d\mu)}$. (In the Fermi liquid theory with s-wave scattering, this product is equal to m/m^* making $\frac{\kappa_e}{\kappa_{WF}}$ exactly equal to unity¹³.) The temperature dependence of the ratio $\frac{\kappa_e}{\kappa_{WF}}$ then describes, in a dimensionless form, the temperature dependence of the specific heat due to the charge-carrying excitations. It is then clear that the prefactor of the asymptotic low-temperature dependence of κ_e should be considered to be anomalously large: The effective bandwidth of the charge-carrying excitations would be of order $0.15K$. Such a small bare scale is essentially impossible.

We are then forced to associate T_0 with a collective scale. Unlike for a bare scale such as a bandwidth, it is entirely reasonable for a collective scale to be this small. On purely phenomenologically grounds, there are two possible interpretations as illustrated in Figs. 1a and 1b.

Distinctive charge-neutral excitations in the normal state: If the charge-carrying excitations indeed have non-fermionic statistics, such that their contribution to the

specific heat has a higher than linear temperature dependence, much of the heat current is necessarily carried by some distinctive charge-neutral excitations. This conclusion is reached as follows. In this picture, the contribution to the thermal conductivity due to the charge-carrying excitations would have to be

$$\frac{\kappa_{charge}}{\kappa_{WF}} \approx \left(\frac{T}{W}\right)^\beta \quad (6)$$

where the exponent $\beta > 0$ reflects both the statistics and the dispersion of the charge-carrying excitations, and W is the effective bandwidth. Since W is expected to be much larger than T_0 , in the measured temperature range κ_{charge} is necessarily much smaller than the measured κ_e as illustrated in Fig. 1a – hence our conclusion. We note that a microscopic theory for such a picture needs to produce a temperature-independent carrier concentration n_{charge} relevant to transport and, at the same time, a specific heat that goes as $T^{1+\beta}$.

Experimentally, the plot¹⁰ of κ_e/T versus T starts to deviate from the low temperature power-law form as the temperature is increased through T_0 . Within our picture, T_0 would be some collective temperature scale associated with a transition that affects mostly the charge-neutral excitations. It follows that the temperature dependence of $\kappa_{neutral}$ is very different from that of κ_{charge} , reflecting the separation of electrons into the charge-carrying and charge-neutral excitations.

Measurement of the heat transport cannot tell us about the quantum number of the charge-neutral excitations. A natural candidate, of course, would be that these are spin-carrying excitations. Measurement of the spin transport would help clarify the nature of these charge-neutral excitations.

Ordering transition in the normal state: An alternative possibility is that, the charge-carrying excitations above T_0 are in fact fermions. A subtle ordering transition takes place around T_0 . Again, it is natural for T_0 to be so small since it is a collective scale.

In this picture, the transition around T_0 leads to a sharp drop in the specific heat but leaves the electrical conductivity largely unaffected. In addition, the fermionic excitations are gapped out at $T \ll T_0$. The reason that κ_e/T goes to zero in the asymptotic low temperature limit is presumably related to what is responsible for the vanishing $\kappa(H=0)/T$. While it is in principle possible, this picture, in its simple form, requires some delicate balance between the quasiparticle contribution and collective contribution to the charge current such that the electrical conductivity is essentially unchanged as the temperature is lowered through T_0 . The collective contribution is of relevance here since disorder is not very strong and pinning is expected to be weak. Whether such a balance can be achieved in specific microscopic models remains to be seen.

Experimentally, $\kappa_e/T\sigma$ at temperatures above T_0 is indeed close to¹⁴, though somewhat larger than, L_0 .

One way to differentiate these two pictures is to study the behavior of κ_e at temperatures high compared to T_0 (but still low enough so that elastic scatterings dominate). In the first picture, the mean free path of the charge-neutral excitations is in general very different from that of the charge-carrying excitations. As illustrated in Fig. 1a, we expect κ_e/κ_{WF} to be very different from unity for $T \gg T_0$. In the second picture, on the other hand, we expect $\kappa_e/\kappa_{WF} \approx 1$ for $T \gg T_0$; see Fig. 1b. Unfortunately, the experiment in PCCO is limited to temperatures not too high compared to T_0 . Analogous experiments in other cuprates may help clarify the situation.

I would like to thank L. Taillefer for sending me a copy of Ref. 10 prior to publication, and E. Abrahams, A. J. Leggett, L. Taillefer, and C. M. Varma for useful discussions. This work was carried out during my participation in the ITP-UCSB program on high temperature superconductivity; I would like to thank the organizers of this program and ITP for hospitality. The work has been supported in part by NSF Grant No. PHY99-07949 (ITP), NSF Grant No. DMR-0090071, and TCSUH.

-
- ¹ P. W. Anderson, *Science* **235**, 1196 (1987).
² *The Theory of Superconductivity in the High- T_c Cuprates* (Princeton University Press, Princeton, NJ 1997).
³ Q. Si, *Phys. Rev. Lett.* **78**, 1767 (1997); *ibid.* **81**, 3191 (1998).
⁴ Q. Si, cond-mat/0003033.
⁵ For a very different proposal to test a class of fractionalized state with topological order, see T. Senthil and M.P.A. Fisher, cond-mat/0006481; see also M. Franz and Z. Tesanovic, cond-mat/0010436.
⁶ V. A. Vas'ko *et al.*, *Phys. Rev. Lett.* **78**, 1134 (1997); *Appl. Phys. Lett.* **73**, 844 (1998).
⁷ Z. W. Dong *et al.*, *Appl. Phys. Lett.* **71**, 1718 (1997).
⁸ N.-C. Yeh *et al.*, *Phys. Rev. B* **60**, 10522 (1999).
⁹ See, e.g., Y. Zhang *et al.*, cond-mat/0001037 (2000).
¹⁰ R. W. Hill, C. Proust, L. Taillefer, P. Fournier, and R. L. Greene, preprint (2000).
¹¹ A. C. Durst and P. A. Lee, *Phys. Rev. B* **62**, 1270 (2000).
¹² Here we consider elastic scatterings only. For inelastic scatterings, there are exceptions to this statement.
¹³ C. Castellani *et al.*, *Phys. Rev. Lett.* **59**, 477 (1987).
¹⁴ This was first emphasized by P. A. Lee.